# University-Based Pulsed Neutron Sources: The LENS Project

David V. Baxter Indiana University

- DVB, J. Cameron (Ph), H. Nann (Ph), E. J. Nelson (GS)
   W. M. Snow (Ph), J. Zwanziger (Ch) (Indiana)
- G. Beaucage (MS-UCi), L. Bronstein (CH-IU), B. Heuser (NE-UIUC), D. Worcester (B-UMo).
- M. Arif (NIST), J. M. Carpenter (ANL), E. B. Iverson (SNS), M. T. Rekveldt (IRI), P. A. Seeger (LANL).

#### **OUTLINE**

- Why build a pulsed neutron source at a University?
- How can it be done?
- What will be done at this facility when it is built?
- Conclusions






**Pulsed** 




#### **Neutron Sources**




#### What is LENS?

- Low Energy Neutron Source: based on lowenergy (p,n) and (p,pn) reactions (E<sub>p</sub><11MeV) in Be.
- The source will be tightly coupled to a cold moderator (solid CH<sub>4</sub> at about 22K).
- LENS will have a variable pulse width (from 100μs to 1.0 ms or more).
- In long-pulse mode, LENS will have a timeaveraged cold neutron intensity greater than the IPNS (1 moderator).
- A small number of scattering instruments will be developed to utilize these neutrons.

## **Existing Infrastructure**



## Low Energy Neutron Source



#### Distinct Features of LENS

- Its low proton energy makes LENS ideal for studies of new moderator materials and designs.
- No high-energy (E>10MeV) neutrons and few γ's allows shorter instrument design, reduced costs, and reduced backgrounds (also UCN candidate).
- Its long (but variable) pulse width makes LENS suitable for Small Angle Neutron Scattering (SANS) and for developing instruments for future long-pulse spallation sources.
- Its low cost (both capital and operating) allows large amounts of beam time to be devoted to education of new users, particularly those in non-traditional fields (Chemistry, Biology)

#### LENS: The numbers

(11MeV, 2.5mA, 22K CH<sub>4</sub> slab moderator)

(5% DF; 900μs proton pulses)

• Primary neutrons: 10<sup>14</sup> /s

Peak Mod. Bright.: 2x10<sup>12</sup> n/cm<sup>2</sup>.s.sr.eV


• Flux @ 5.0m: **7x10**<sup>6</sup> n/cm<sup>2</sup>.s

(0.7-33meV, uncol.)

• SANS flux: >3x10<sup>5</sup> n/cm<sup>2</sup>.s (1.06-0.16nm,col.)

Based on MCNP and NISP calculations. Does not include: Grooved moderator, composite target and/or moderator, ...

## Neutron Energy Distribution



Derived from MCNP calculations similar to those vetted on the IPNS-H moderator (Iverson99).


## So why does LENS work?

#### **Feature**

#### **LENS/IPNS**


<ul> <li>Nuclear efficiency</li> </ul>	1/2500
<ul> <li>Larger average current</li> </ul>	200
<ul> <li>No decouplers/poisons</li> </ul>	6
<ul> <li>Geometric Coupling</li> </ul>	2-5
<ul> <li>Lower energy spectrum</li> </ul>	1.2-5
<ul> <li>Overall</li> </ul>	1-3

### **Conventional SANS**



Roughly equivalent to the **SAD** instrument at **IPNS** 

## Spin-Echo SANS



- Encode momentum info in neutron SPIN!
- •Resolution is decoupled from the incident beam phase space! (NO COLLIMATION NEEDED in the encoded direction!)
- •Max length scale probed is set by the maximum B field; 0.1T → microns!
- •Variations on the theme allow high-resolution diffraction, SAS, reflectometry etc. (stay around for Roger Pynn's talk!)

## **CONCLUSIONS**

- LENS will have a time-averaged cold neutron intensity greater than that available from a single moderator at the IPNS in a university setting!
- The LENS time structure is most suited to SANS (for max. flux) but is flexible for instrumentation and moderator development and can be used for a variety of other techniques with suitable instrumentation.
- LENS can play an important role in building the user base needed to support the SNS, it is our hope that a network of such sources will be built.
- By combining SANS, SESANS, and phase-contrast radiography, LENS will be able to investigate materials structures over 7 orders of magnitude in length scale.